

Allelochemical Interactions Between Sponges and Corals

JAMES W. PORTER¹ AND NANCY M. TARGETT²

¹Zoology Department, University of Georgia, Athens, Georgia 30602, and ²College of Marine Studies, University of Delaware, Lewes, Delaware 19958

Abstract. The existence of chemical-biological interactions is routinely invoked to explain patterns of coexistence between neighboring organisms. This study characterizes the consequences of these interactions in interspecific space competition between the neighboring species *Plakortis halichondroides*, the liver sponge, and *Agaricia lamarcki*, the sheet coral. This sponge/coral association was studied *in situ* both at points of natural contact and following manipulations that artificially brought the sponge and coral together. *Plakortis* kills *Agaricia* upon direct contact and upon indirect contact (*i.e.*, waterborne metabolites only). *Plakortis* creates a dead zone of coral around its base as it overgrows the coral. The effect of either direct or indirect contact by *Plakortis* is to reduce: (1) the number of zooxanthellae in *Agaricia*, (2) the weight of chlorophyll *a* per unit area of the coral, and (3) the weight of tissue nitrogen per unit area of the coral.

The necrotic effect also evidences itself as changes in oxygen flux characteristics such as significant increases in the compensation point and the nocturnal respiration rate, and significant reductions in the maximum net and gross photosynthetic rates. As a consequence, the diel integrated production to respiration ratio falls below unity for *Agaricia* colonies in contact with *Plakortis*; this does not occur for coral without neighboring sponges.

Because direct contact between *Plakortis* and *Agaricia* is not necessary to effect stress in the coral, the presence of active chemical metabolites from *Plakortis* is suggested. Thus, mechanical abrasion is excluded as the sole mechanism of dominance by the sponge.

Introduction

Sessile and sedentary coral reef organisms frequently compete for space and food, adopting mechanisms to

minimize fouling or overgrowth by epibionts and maximize their own space-capture abilities. Several biological mechanisms that mediate ecologically significant interactions among coral reef organisms have been described. For example, scleractinian corals effect extracoelenteric damage to neighbors via extended mesenterial filaments and long sweeper tentacles (Francis, 1973; Lang, 1973; Richardson *et al.*, 1979; Wellington, 1980; Sheppard, 1982). Likewise, hydrocorals and octocorals can move onto and spread across scleractinians, and thereby compete successfully for space with reef-building corals (Wahle, 1980; La Barre and Coll, 1982; Tursch and Tursch, 1982). Bryozoans employ sweeper appendages which are effective in competition and in prevention of fouling (Jackson, 1977). These structures are used in specific behaviors that involve the recognition of potential competitors and the direction of interference mechanisms against them.

Chemical defense mechanisms have also been suggested. These mechanisms have demonstrable effects on species distributions and individual survivorship in terrestrial plant communities (Fraenkel, 1969; Whittaker and Feeny, 1971; Rosenthal and Janzen, 1979; Meinwald, 1982; Targett and Isman, 1986), and have recently been reviewed in ecological contexts for marine communities (Barbier, 1981; Bak *et al.*, 1982; Fenical, 1982; Norris and Fenical, 1982; Palumbi and Jackson, 1982; Colwell, 1983; Faulkner and Ghiselin, 1983; Steinberg, 1984; Scheuer, 1985; Bakus *et al.*, 1986).

Unusual secondary metabolites have been isolated from numerous sessile solitary and colonial coral reef organisms (Tursch *et al.*, 1978; Cimino *et al.*, 1983; Sullivan *et al.*, 1983; Coval *et al.*, 1984; Bandurraga and Fenical, 1985; Kashman *et al.*, 1985; and Coll *et al.*, 1985). Animals that contain unusual secondary metabolites of-

ten prove to be toxic in bioassay examinations (Bakus and Thun, 1979; Bakus, 1981; Coll *et al.*, 1982a, 1983; Targett *et al.*, 1983; Gerhart, 1984; McCaffrey and Eidean, 1985; Thompson, 1985; Thompson *et al.*, 1985; LaBarre *et al.*, 1986a, b). In addition to their role in organism defense, secondary metabolites are also implicated in the maintenance living space (Jackson and Buss, 1975; Green, 1977; Jackson, 1977; Sheppard, 1979, 1982; Sammarco *et al.*, 1983).

Sponges are remarkable because they lack specialized organs and behaviors, and yet are successful in an environment where such adaptations are common. However, sponges do contain a variety of bioactive secondary metabolites. More than three dozen compounds with lethal or growth inhibitory properties are described from tropical sponges in reviews by Russell and Saunders (1967), Sigel *et al.* (1969), Martin and Padilla (1973), Baker and Murphy (1976), Hollenbeak *et al.* (1976), Faulkner (1977), Cimino (1977), Minale (1978), Kaul and Sinderman (1978), and Hashimoto (1979). These researchers characterize the compounds chemically and occasionally list effects on organisms of direct interest to man, but rarely include data on the ecological importance of the compound or demonstrate effects on other marine organisms likely to have frequent encounters with the species. Recent work continues to identify unusual secondary metabolites from sponges (*e.g.*, Fusetani *et al.*, 1981; Tachibana *et al.*, 1981; Carmely *et al.*, 1983; Cimino *et al.*, 1983; Gonzalez *et al.*, 1984; Nakatsu *et al.*, 1983, 1984; Manes *et al.*, 1985; Braekman *et al.*, 1985; Walker *et al.*, 1985; Nakamura *et al.*, 1986; Mayol *et al.*, 1986) and also identifies several ecological contexts in which the metabolites might function (Thompson *et al.*, 1982; Cimino *et al.*, 1982).

Sullivan *et al.* (1983) demonstrate that the burrowing sponge *Siphonodictyon* sp. secretes a guanidine-containing sesquiterpene, siphonodictidine, in its mucus. This compound kills adjacent coral tissue, thereby preventing the coral from overgrowing the sponge's oscular chimneys. The compound stimulates coral respiration and it has been speculated that increased respiration rate and decreased photosynthetic rate would probably result in death for the coral. However, Sullivan *et al.* (1983) point out that because so many factors can affect hard corals under aquarium conditions, one must be cautious in extrapolating to the long-term effects of siphonodictidine at subacute concentrations. Environmental information on the effects of these kinds of compounds *in situ* is needed.

In this paper, we characterize the consequences of chemical-biological interactions in interspecific space competition between the coral *Agaricia lamarcki* (Milne Edwards & Haime, 1848) and the sponge *Plakortis halichondroides* (Wilson, 1902). Interactions between *Pla-*



Figure 1. Natural associations of the sponge *Plakortis halichondroides* and the coral *Agaricia lamarcki* show a bleached area at the zone of contact. The necrotic area is evident in this photograph on the left side of the coral along the region adjacent to the sponge and above the 1.0 cm scale bar in the center. (Salt River Canyon, St. Croix, 25 m depth; photograph by R. S. Smith).

kortis and *Agaricia* are common in St. Croix, U. S. Virgin Islands, and in Jamaica where *Plakortis* overgrows living coral, utilizing the newly dead coral skeletons as a primary point of attachment and growth (N. Targett, J. Neigel, J. Porter, unpub. data). Upon physical contact or proximity of less than five centimeters to the sponge, *Agaricia* bleaches and shows a marked necrosis in the region of contact (Fig. 1).

Using the *Plakortis/Agaricia* interaction as a basis for study, we: (1) describe the *in situ* effect of *Plakortis halichondroides* on *Agaricia lamarcki* both at points of natural contact and following manipulations that artificially bring together the sponge and coral, (2) quantify the *in situ* effects of direct sponge contact and indirect sponge contact (*i.e.*, exposure to whole sponge exudates only)

on photosynthetic and respiratory oxygen fluxes in the coral, and (3) suggest the long-term effects on tissue biomass, growth rate and survival of *Agaricia* colonies growing in association with *Plakortis* sponges.

Materials and Methods

Site description

This research was conducted utilizing the NOAA underwater habitat, Hydrolab, located at a depth of 17 m on the north coast of St. Croix, U. S. Virgin Islands. Experimental material and line transect data were collected from the east coral reef slope of the Salt River Canyon within the research area available from the habitat.

Field studies

Transects ten meters in length and two meters wide (for a total of 20 m² in each transect area) were layed parallel to depth contours at 20 and 30 m. The number of *Plakortis halichondroides* colonies within one meter of the line were counted. The number of times *P. halichondroides* grew within five centimeters of living tissue of *Agaricia lamarcki* was also noted. Additionally, a swimming census covering 1000 m² was conducted between the 20 and 30 m depth contours to record all species of scleractinian corals adjacent to *P. halichondroides* colonies.

Biomass determinations

Coral tissue was removed from the surface of the coral skeleton using a Water Pik (Johannes and Wiebe, 1970) with filtered seawater. The tissue slurry was homogenized for one minute in a blender and triplicate aliquots were removed for analysis of nitrogen content, zooxanthellae cell density, and chlorophyll *a* concentration. Standard protocols for sample preparation and analysis were used for fluorometric determination of chlorophyll *a* (Holm-Hansen *et al.*, 1965), hemocytometric counts of zooxanthellae (Muscatine *et al.*, 1984), and micro-Kjeldahl nitrogen digestions (Parsons *et al.*, 1984) as described for coral tissue analysis in Porter *et al.* (1984). Coral surface area was determined for *Agaricia lamarcki* using Marsh's (1970) aluminum foil overlay technique.

Oxygen flux determinations

Oxygen flux in three experimental chambers and ambient PAR light flux on a 4- π quantum sensor were recorded on a Datel magnetic tape data logger at four minute intervals during the 24-hour experimental incubations. Tapes from each experiment were read into a Tektronics 4054 microcomputer where consecutive readings were subtracted from one another. The result

was then multiplied by the volume of water in the incubation chambers to give net rates of oxygen production or consumption. These oxygen flux values were normalized to biomass units of surface area, and mass of nitrogen and chlorophyll *a*.

Coral pieces were cut from larger colony rosettes and were used in physiological experiments after they had acclimated for 24 hours. Experimental corals were monitored prior to treatment as well as after treatment, and therefore each served as its own pre-treatment control.

Photosynthesis *versus* irradiance curves were constructed for each control or experimental condition by computing average fluxes over small intervals of irradiance (maximum interval 50 $\mu\text{Em}^{-2} \text{s}^{-1}$). These points were fitted to a hyperbolic tangent function (Jassby and Platt, 1976) which represented the best-fit for patterns of coral productivity (Porter, 1980; Chalker, 1980). The program derived from this function allowed accurate statistical determinations of (1) α the initial slope of the photosynthesis-irradiance curve, (2) I_c , the light compensation point, (3) I_k , the break point, where the P:I curve approached p_{max} , (4) p_c net max and p_c gross max, the maximum net and gross coral head production, (5) r_c , coral nocturnal respiration rate, and (6) an integrated, diel P/R ratio. We have fully described the interpretive model for coral photosynthesis/respiration ratios elsewhere (Muscatine *et al.*, 1981; Muscatine *et al.*, 1984; Porter, 1985).

Experimental design

Eleven null hypotheses were formulated to investigate the consequences to *Agaricia lamarcki* of association with *Plakortis halichondroides*. These null hypotheses included information on biomass patterns (9–11), physiological rates (1–6), and carbon budget totals (7–8) for corals before and after contact with sponges (direct contact) or with sponge exudates (indirect contact). The following null hypotheses were advanced:

1. α Before = α After Exposure
2. I_k Before = I_k After Exposure
3. I_c Before = I_c After Exposure
4. p_c net max Before = p_c net max After
5. p_c gross max Before = p_c gross max After
6. r_c night Before = r_c night After
7. P/R Ratio Before = P/R Ratio After
8. P/R Ratio Exceeds 1.0 Before; Less Than 1.0 After
9. [Zooxanthellae] Before = [Zooxanthellae] After
10. [Pigment] Before = [Pigment] After
11. [Nitrogen] Before = [Nitrogen] After

The four experimental treatments listed below were designed to test these hypotheses, *i.e.*, they were designed to reject the contention that there were no significant

effects on coral biomass or oxygen flux patterns after contact with sponges. The *in situ* manipulations included:

1. Control, no contact or history of contact between *Plakortis halichondroides* and the examined *Agaricia lamarcki* colonies.

2. Indirect contact, *A. lamarcki* colonies in contact with seawater containing exudates from coarsely mashed *P. halichondroides*.

3. Direct contact, *A. lamarcki* colonies in physical contact with pieces of *P. halichondroides* cut from the reef and tied to the coral colonies for 24 hours prior to the oxygen flux measurements.

4. Direct contact, *A. lamarcki* colonies naturally occurring adjacent to uninjured *P. halichondroides* colonies on the reef.

Experimental condition (4) was of greatest ecological interest since, of the three experimental treatments, it most closely recorded conditions of natural contact on the reef.

Oxygen consumption by the electrodes (all experiments) and the injected sponge water (Experiment 2 only) were determined just prior to the experimental incubations. These chemical oxygen demands were added to oxygen fluxes inside the chambers to give the metabolic activity of the coral alone.

The injured and uninjured sponges used in the last two treatments (3 and 4, respectively) were removed from the coral's surface prior to oxygen flux measurements on the coral colony. Since sponge metabolic rates *per se* were not of direct interest in the experimental design, and since Reisinger (1971) has shown that small chambers are unsuitable for the determination of oxygen flux characteristics for sponges like *Plakortis* with high pumping rates, the metabolic activity of *Plakortis* was not determined.

As a control on the experimental method employed, the effects of contact with sponges such as *Agelas*, *Haliclona*, and *Verongula*, which did not cause bleaching and necrosis, were also monitored. Finally, to examine metabolites released from *P. halichondroides*, an *in situ* chemical sampling pump concentrated organic compounds from the water surrounding uninjured colonies of this species.

Results

Field studies

All coral species found in the transect area are overgrown and killed by *Plakortis halichondroides* (Table I). These fourteen species are from eight of the nine Caribbean coral families (the remaining family, the Acroporidae, was not found in this reef zone). The overgrown corals

Table I

Hermatypic scleractinian corals killed in situ by Plakortis halichondroides (Wilson, 1902)

| | |
|-------|---|
| I. | Family Astrocoeniidae Koby, 1890 |
| | 1. <i>Stephanocoenia michelinii</i> Milne Edwards & Haime, 1848 |
| II. | Family Pocilloporidae Gray, 1842 |
| | 2. <i>Madracis decactis</i> (Lyman, 1859) |
| III. | Family Agariciidae Gray, 1847 |
| | 3. <i>Agaricia agaricites</i> (Linnaeus, 1758) |
| | 4. <i>Agaricia lamarcki</i> Milne Edwards & Haime, 1851 |
| IV. | Family Siderastreae Vaughan & Wells, 1943 |
| | 5. <i>Siderastrea siderca</i> (Ellis & Solander, 1786) |
| V. | Family Poritidae Gray, 1842 |
| | 6. <i>Porites astreoides</i> Lamarck, 1816 |
| VI. | Family Faviidae Gregory, 1900 |
| | 7. <i>Diploria labyrinthiformis</i> (Linnaeus, 1758) |
| | 8. <i>Diploria strigosa</i> (Dana, 1846) |
| | 9. <i>Colpophyllia natans</i> (Houttuyn, 1772) |
| | 10. <i>Montastraea annularis</i> (Ellis & Solander, 1786) |
| | 11. <i>Montastraea cavernosa</i> (Linnaeus, 1766) |
| VII. | Family Meandrinidae Gray, 1847 |
| | 12. <i>Meandrina meandrites</i> (Linnaeus, 1758) |
| VIII. | Family Mussidae Ortmann, 1890 |
| | 13. <i>Mycetophyllia lamarckiana</i> Milne Edwards & Haime, 1848-1849 |
| | 14. <i>Mycetophyllia ferox</i> Wells, 1973 |

als are more similar morphologically than taxonomically: they are all horizontally flattened, and as such, provide a level substratum on which *Plakortis* can grow. The only exception to this pattern is the finger coral, *Madracis mirabilis*, which is killed at the base of its branches in contact with the sponge.

Approximately one-third ($34.0 \pm 6.8\%$) of all *Plakortis* colonies occur on or directly adjacent to living coral (Table II). The remaining members of the population grow on stable substrata, many of which are recently dead coral plates. Almost half of the corals with sponges on or near them show signs of bleaching and necrosis ($40.8 \pm 3.4\%$, Table II), but all coral specimens show tissue death in the area directly underneath the sponge.

Biomass

Our results clearly demonstrate a marked effect on algal densities, algal pigment concentration, and coral tissue mass for all of the experimental conditions relative to the control condition (Table III). The number of zooxanthellae per unit area, and as a consequence, the chlorophyll *a* per unit area of coral, decreases by a factor of three after contact with injured sponges, uninjured sponges, and sponge water (Table III). The amount of chlorophyll *a* per algal cell does not follow a consistent pattern under the experimental treatments; only after exposure to sponge water does the mass of chlorophyll *a* per algal cell decline significantly (Table III).

Table II

Plakortis halichondroides population density and interactions with scleractinian corals ($\bar{x} \pm$ one S.D., $n = 4$)

| Station name | Total number of sponge colonies per 20 m ² | Total number of sponge colonies within 5 cm of coral | % of sponge population in contact with coral | Number of coral colonies within 5 cm of sponges | Number of coral colonies showing visible signs of stress | % of contacted corals showing signs of stress |
|--------------|---|--|--|---|--|---|
| 20m Right | 62 | 18 | 29 | 16 | 6 | 38 |
| 30m Left | 39 | 13 | 33 | 13 | 5 | 38 |
| 30m Right | 105 | 32 | 30 | 33 | 15 | 45 |
| 20m Left | 90 | 40 | 44 | 36 | 15 | 42 |
| | 74 ± 29 | 26 ± 12 | 34.0 ± 6.9 | 24 ± 12 | 10 ± 5 | 40.8 ± 3.4 |

Nitrogen mass per unit area of coral tissue decreases by a factor of two (Table III). This effect cannot be explained solely by the loss of zooxanthellae since they constitute only 7% of the coral tissue biomass (Porter and Muscatine, in prep.). Coral tissue lysis also occurs (Fig. 2). The effect of these combined plant and animal tissue responses is to create a bleached zone, or necrotic halo, on the coral in the vicinity of the sponge that is visually obvious from a meter away (Fig. 1).

Oxygen flux

Contact with *Plakortis* effects both oxygen production and oxygen consumption (Fig. 3). Oxygen consumption more than doubles for corals in direct contact with injured or uninjured sponges, rising from approximately 6 to 12 $\mu\text{gO}_2\text{cm}^{-2}\text{h}^{-1}$ (Table IV). Coral respiration rate stays the same during injection of sponge water in the short-term indirect contact experiments. However, the maximum net photosynthetic rate drops by almost half for both direct contact with injured or uninjured sponges and for indirect contact with sponge water. These lowered photosynthetic rates occur within eight minutes of injecting sponge water, suggesting that sponge metabolites rapidly diminish this coral species' photosynthetic

capacity. Rising respiration rates and diminishing production rates tend to offset one another mathematically. And hence, only for indirect contact with sponge water, where production falls but respiration does not rise concomitantly, does the maximum gross production show a significant decline.

The compensation light intensity is significantly higher under all experimental treatments relative to the control (Fig. 3; Table IV). This demonstrates that more light is needed for corals near or adjacent to sponges to meet their basal metabolic demands through photosynthesis than for corals at a distance from sponges. Further, it suggests that coral production balances coral respiration later in the morning and stops earlier in the afternoon for corals near sponges (Fig. 3).

Given the lowered production rates and increased respiration rates observed under the experimental treatments, it is not surprising that coral P/R ratios are also substantially lower among corals in contact with sponges (Table IV). The overall effect is that while the integrated P/R ratio is always at or above unity in control corals, it is always below 1.0 in corals exposed to injured or uninjured sponges under field conditions (Table IV). P/R ratios below 1.0 are never found for this species in Jamaica,

Table III

Biomass variation ($\bar{x} \pm$ one S.D., $N = 3$) for colonies of the sheet-coral, *Agaricia lamarcki* under different exposures to the liver sponge, *Plakortis halichondroides* (see Figs. 1 & 2)

| Characteristic | Units | Experimental treatment | | | |
|----------------|------------------------------------|------------------------|---------------------------|-----------------------------|-------------------------------|
| | | No contact (Control) | Contact with sponge water | Contact with injured sponge | Contact with uninjured sponge |
| Zooxanthellae | 10^6 cells cm^{-2} | 1.04 ± 0.08 | 0.36 ± 0.08* | 0.38 ± 0.06* | 0.40 ± 0.14* |
| Pigments | pg Chl <i>a</i> cell ⁻¹ | 10.02 ± 0.10 | 5.82 ± 0.93* | 9.81 ± 4.05 | 10.57 ± 1.88 |
| Pigments | $\mu\text{g Chl a cm}^{-2}$ | 10.39 ± 0.86 | 2.08 ± 0.13* | 3.79 ± 1.61* | 2.84 ± 1.00* |
| Nitrogen | mg TN cm^{-2} | 0.57 ± 0.06 | 0.25 ± 0.03* | 0.25 ± 0.16 | 0.25 ± 0.10* |

* Means significantly different from the Control ($P \leq 0.05$, ANOVA).




Figure 2. After two hours of experimentally induced contact between *Agaricia lamarcki* and *Plakortis halichondroides* (a, top), a bleached, necrotic area appears on the coral (b, bottom). (Salt River Canyon, St. Croix, 25 m depth; photograph by N. M. Targett).

even over greater depth ranges (Porter and Muscatine, in prep.), and therefore these values for coral colonies in contact with sponges are indicative of unsustainable carbon deficits.

In all cases, the differences observed between control and experimental coral oxygen flux patterns demonstrate significantly reduced photosynthetic capacity in coral colonies in contact with sponges or their metabolites. Therefore, the null hypotheses (no discernable effect) must be rejected for hypotheses (3) compensation point, (4) net production, (5) gross production, (6) nocturnal respiration rate, and (7) integrated P/R ratio under either field conditions or idealized "cloudless day" illumination. Hypothesis (8) P/R ratio > 1.0 is also falsified.

Other species of sponges (*Agelas conifera*, *Haliciona rubens*, and *Verongula* sp.) were also placed in contact with *Agaricia lamarcki*. They did not bleach the coral, thus indicating that mechanical irritation and pressure

are not responsible for the effect observed with *P. halichondroides*.

Crude organic extracts were isolated from *Plakortis halichondroides* and coated onto synthetic cellulose pads, "tuffy sponges." When tied to living coral, these extract-soaked pads caused bleaching within 24 hours. Control pads (uncoated or coated with ether solvent only) produce no effect. While the exact nature of these organic compounds is still unknown, comparative thin layer chromatography of crude extracts from whole *P. halichondroides* and compounds isolated from waters surrounding uninjured *Plakortis* suggests that compounds in the surrounding water are the same as those produced naturally by the sponge.

Discussion

The sponge *Plakortis halichondroides* actively inhibits the metabolism and tissue survival of adjacent corals. Inhibition results from both direct and indirect contact (waterborne metabolites only) suggest that *Plakortis* uses allelochemicals as one means to secure and occupy space on the reef.

Plakortis halichondroides bleaches *Agaricia lamarcki* and causes marked tissue necrosis. The effects on coral biomass and coloration are sufficiently dramatic that it is possible to survey these interactions visually from some distance above the reef surface. The loss of zooxanthellae from corals following exposure to sponges or sponge exudates parallels the loss of symbiotic algae during other natural stresses such as abnormal temperatures, salinity fluctuations, or high rates of sedimentation (Porter, 1987).

These coral biomass reductions contribute to the profound effects that sponges have on coral oxygen metabolism. Paralleling the loss of zooxanthellae is a significant decline in primary production. Although the algae that remain appear to have normal concentrations of photosynthetic pigments, the few remaining zooxanthellae cannot compensate for the overall loss of algae. Further, despite the fact that there is significantly less coral tissue per unit area on corals adjacent to sponges than on corals without sponge contact, the respiration rate is still significantly higher. This respiratory increase suggests a stress or repair-metabolism response to the active sponge metabolites.

The overall effect on the P/R formula of decreasing the numerator and increasing the denominator is to lower the ratio below one for *Agaricia lamarcki*. These suboptimal values are not found in this photoautotrophic coral species. For example, even to depths of 50 m, *Agaricia* has an annual integrated P/R ratio of 1.13 (Porter and Muscatine, in prep.).

The ecological role of specific secondary metabolites

Table IV

Variation ($\bar{x} \pm 95\%$ confidence limits, $n = 6$) in photosynthesis-light utilization characteristics for in situ colonies of *Agaricia lamarcki* under different exposure to *Plakortis halichondroides* (see Fig. 3)

| Characteristic | Units | Experimental treatment | | | |
|----------------------------|--|------------------------|---------------------------|-----------------------------|-------------------------------|
| | | No contact (Control) | Contact with sponge water | Contact with injured sponge | Contact with uninjured sponge |
| α (cm^2) | $\mu\text{gO}_2 \text{ cm}^{-2} \text{ h}^{-1} \mu\text{E}^{-1} \text{ m}^{-2} \text{ s}^{-1}$ | 0.222 ± 0.046 | 0.140 ± 0.058 | 0.266 ± 0.141 | 0.303 ± 0.196 |
| I_k (cm^2) | $\mu\text{Em}^{-2} \text{ s}^{-1}$ | 137.82 ± 9.16 | 129.64 ± 27.93 | 109.40 ± 25.10 | 101.32 ± 36.12 |
| I_c (cm^2) | $\mu\text{Em}^{-2} \text{ s}^{-1}$ | 27.69 ± 1.70 | $60.54 \pm 15.58^*$ | $76.90 \pm 21.00^*$ | $85.84 \pm 19.68^*$ |
| p_c net max | $\mu\text{gO}_2 \text{ cm}^{-2} \text{ h}^{-1}$ | 24.88 ± 1.95 | $10.64 \pm 2.22^*$ | $17.90 \pm 4.19^*$ | $15.05 \pm 7.12^*$ |
| p_c gross max | $\mu\text{gO}_2 \text{ cm}^{-2} \text{ h}^{-1}$ | 30.66 ± 1.95 | $14.78 \pm 2.22^*$ | 29.16 ± 4.19 | 30.98 ± 7.12 |
| Average r_c night | $\mu\text{gO}_2 \text{ cm}^{-2} \text{ h}^{-1}$ | 5.77 ± 0.21 | 4.14 ± 0.29 | $11.26 \pm 0.80^*$ | $15.93 \pm 1.09^*$ |
| P_c gross/ R_c 24 h | Ratio (field) | 1.98 ± 0.51 | 1.37 ± 0.31 | $0.83 \pm 0.34^*$ | $0.73 \pm 0.23^*$ |

* Means significantly different from the Control ($P \leq 0.05$).

in marine organisms is known in only a few cases (Webb and Coll, 1983; Sullivan *et al.*, 1983; Morse and Morse, 1984; Targett *et al.*, 1986; Pawlik, 1986). In the boring sponge *Siphonodictyon*, mucus-borne metabolites kill corals (Sullivan *et al.*, 1983). Our study demonstrates that chemical-biological interactions are important in non-boring sponge species and in over-growth, not just anti-fouling, processes.

Several toxic exudates from soft corals appear to be




Figure 3. Net photosynthesis-irradiance curves are graphed for *Agaricia lamarcki* colonies ($n = 3$) from 20 m depth under different levels of contact with *Plakortis halichondroides*. The curves are fitted to a hyperbolic tangent function (see Table IV for the statistical comparisons between curves). Experimental Condition 1: control, no contact with *Plakortis* (open circles). Experimental Condition 2: indirect contact with water containing exudate from coarsely mashed *Plakortis* (solid squares). Experimental Condition 3: direct contact with injured *Plakortis* (solid triangles) (see Fig. 2). Experimental Condition 4: direct contact with uninjured *Plakortis* (solid diamonds) (see Fig. 1). Experimental Condition 4 most closely mimics conditions of natural contact on the reef.

responsible for causing localized mortality in hard corals, since extensive mortality of hard corals can occur even when direct contact is not established with the soft coral (Sammarco *et al.*, 1985). Coll *et al.* (1982b) isolated secondary metabolites from the water surrounding two species of octocorals that were identical to those isolated from the octocorals themselves. Flexibilide, dihydroflexibilide, sarcophine, and sarcophytoxide were isolated from crude extracts of *Simularia flexibilis* and *Sarcophyton crassocaule*, respectively, and from their surrounding water when sampled *in situ* (Coll *et al.*, 1982b). Flexibilide and sarcophytoxide cause death in scleractinian corals at concentrations > 10 ppm (Webb and Coll, 1983). These compounds are also structurally related to simulariolide, a potent algal growth inhibitor (Tursch, 1976).

The sponge *Plakortis halichondroides* contains numerous unusual secondary metabolites such as cyclic peroxides and aromatic lactones (Faulkner *et al.*, 1979; Stierle and Faulkner, 1980). It is possible that the cyclic peroxides cause the bleaching response described above and that the phenolic compounds contribute to tissue lysis. The closely related congener, *Plakortis zygompha*, contains (Z)-7-methyl-4-octen-3-one and several derivatives of 3-hydroxy-4-hydroxymethyl-4-pentenoic acid (Faulkner and Ravi, 1980). These may be involved in both the coral biomass changes and the oxygen flux modifications observed for the *Plakortis/Agaricia* association.

Between 30 and 40 percent of all *Plakortis* colonies on the reef are contiguous with scleractinian corals. Of these contacts, half show bleaching and necrosis on the coral tissue near the sponge; all show death of the coral tissue underneath the sponge. More importantly, based on the results of our *in situ* manipulations, the same effect is observable prior to the establishment of direct sponge-

coral contact. This implies that the sponge exudes waterborne metabolites which have a detrimental effect on coral respiration and photosynthesis. If even naturally low concentrations of these waterborne metabolites are effective at suppressing coral photosynthesis, then there is the distinct possibility that sponge exudates may influence ecosystem productivity in the deeper zones of the reef.

Plakortis kills and colonizes the dead skeletons of at least 14 scleractinian coral species. These species are distributed in all 8 Caribbean coral families located in the 20–30 m survey area. Our demonstration that toxins from *Plakortis* are so effective against corals, the otherwise most successful order of benthic invertebrates on the reef, suggests that further investigation of this interaction is warranted. Future research should focus on the mechanisms of action and chemical structure of the compounds that inhibit photosynthesis, stimulate respiration, and reduce coral colony biomass. The range of potency and the half-life of these allelochemicals once released into the water column is also of critical interest. Finally, information on the long-term survival of corals at varying distances from sponges, and the sources of mortality for *Plakortis* would be of great value to our understanding of the chemical ecology of this association. Allelochemistry gives *Plakortis* a measurable advantage in space competition with scleractinian corals, but is only one factor among many other biotic and abiotic factors which determine its population structure and dynamics.

Acknowledgments

We thank the Hydrolab staff for their support during Missions 84-10 and 85-4, and particularly thank aquanauts George Schmahl, Joe Schubauer, Paul Preston from 84-10, and Nick Vrolijk, Ray Jakubczak, and Robbie Smith from 85-4. We are also grateful to topside scientists Hany Salam (84-10 and 85-4) and Susan Sennett (85-4). Dr. Willard D. Hartman identified our *Plakortis* specimens. This work is supported by grants from the Hydrolab Program of NOAA and Office of Naval Research Grant Number N-000-14-84-K-0571.

Literature Cited

- Bak, R. P. M., R. M. Termaat, and R. Dekker. 1982. The complexity of coral interactions: influence of time, location of interaction, and epifauna. *Mar. Biol.* **69**: 215–222.
- Baker, J. T., and V. Murphy. 1976. *Compounds from Marine Organisms. Volume 1* CRC Press, Cleveland. 226 pp.
- Bakus, G. J. 1981. Chemical defence mechanisms and fish feeding behavior on the Great Barrier Reef, Australia. *Science* **211**: 497–499.
- Bakus, G. J., N. M. Targett, and B. Schulte. 1986. Chemical ecology of marine organisms: an overview. *J. Chem. Ecol.* **12**: 951–987.
- Bakus, G. J., and M. Thun. 1979. Bioassays on the toxicity of Caribbean sponges. *Colloq. Internat. C. N. R. S.* **291**: 417–422.
- Bandurraga, M. M., and W. Fenical. 1985. Isolation of the muricins: evidence of a chemical adaptation against fouling in the marine octocoral *Muricea fructicosa* (Gorgonacea). *Tetrahedron* **41**: 1057–1065.
- Barbier, M. 1981. Marine chemical ecology: the roles of chemical communication and chemical pollution. Pp. 148–186 in *Marine Natural Products. Chemical and Biological Perspectives Vol. IV*, P. J. Scheuer, ed. Academic Press, New York.
- Braekman, J. C., D. Daloz, M. Kaisin, and B. Moussiaux. 1985. Ichthyotoxin sesterperenoids from New Guinean sponge *Carterospongia foliasens*. *Tetrahedron* **41**: 4603–4614.
- Carmely, S., Y. Loya, and Y. Kashman. 1983. Siphonellinol, a new triterpene from the marine sponge *Siphonochalina siphonella*. *Tetrahedron Lett.* **24**: 3673–3676.
- Chalker, B. E. 1980. Modelling light saturation curves for photosynthesis: an exponential function. *J. Theoret. Biol.* **84**: 205–215.
- Cimino, G. 1977. A survey of sesquiterpenoids from marine sponges. Pp. 61–86 in *Marine Natural Products Chemistry*, D. J. Faulkner and W. Fenical, eds. Plenum Press, New York.
- Cimino, G., S. deRosa, S. deStefano, R. Self, and G. Sodano. 1983. The bromocompounds of the true sponge *Verongia aerephoba*. *Tetrahedron Lett.* **24**: 3029–3032.
- Cimino, G., S. deRosa, S. deStefano, and G. Sodano. 1982. The chemical defense of four Mediterranean nudibranchs. *Comp. Biochem. Physiol.* **73B**: 471–474.
- Coll, J. C., B. F. Bowden, D. M. Tapiolas, and W. C. Dunlap. 1982a. *In situ* isolation of allelochemicals released from soft corals (Coelenterata: Octocorallia): a totally submersible sampling apparatus. *J. Exp. Mar. Biol. Ecol.* **60**: 293–299.
- Coll, J. C., B. F. Bowden, D. M. Tapiolas, R. H. Willis, P. Djura, M. Streamer, and L. Trott. 1985. Studies of Australian soft corals-XXXV: the terpenoid chemistry of soft corals and its implications. *Tetrahedron* **41**: 1085–1092.
- Coll, J. C., S. LaBarre, P. W. Sammarco, W. T. Williams, and G. J. Bakus. 1982b. Chemical defenses in soft corals (Coelenterata: Octocorallia) of the Great Barrier Reef: a study of comparative toxicities. *Mar. Ecol. Progr. Ser.* **8**: 271–278.
- Coll, J. C., D. M. Tapiolas, B. F. Bowden, L. Webb, and H. Marsh. 1983. Transformation of soft coral (Coelenterata: Octocorallia) terpenes by *Ovula ovum* (Mollusca: Prosobranchia). *Mar. Biol.* **74**: 35–40.
- Colwell, R. R. 1983. Biotechnology in the marine sciences. *Science* **222**: 19–24.
- Coval, S. J., P. J. Scheuer, G. K. Matsumoto, and J. Clardy. 1984. Two new xenicin diterpenoids from the octocoral *Anthelia edmondsoni*. *Tetrahedron* **40**: 3823–3828.
- Faulkner, D. J. 1977. Interesting aspects of marine natural products chemistry. *Tetrahedron* **33**: 1421–1443.
- Faulkner, D. J., R. W. Armstrong, P. Djura, M. D. Higgs, B. N. Ravi, D. B. Stierle, and S. J. Wratten. 1979. Some chemical constituents in Caribbean sponges. *Colloq. Internat. C. N. R. S.* **291**: 401–406.
- Faulkner, D. J., and M. T. Ghiselin. 1983. Chemical defense and evolutionary ecology of dorid nudibranchs and some other opisthobranch gastropods. *Mar. Ecol. Progr. Ser.* **13**: 295–301.
- Faulkner, D. J., and B. N. Ravi. 1980. Metabolites of the marine sponge *Plakortis zygompha*. *Tetrahedron Lett.* **21**: 23–26.
- Fenical, W. 1982. Natural products chemistry in the marine environment. *Science* **215**: 923–928.
- Fraenkel, G. 1969. Evolution of our thoughts on secondary plant substances. *Etamol. Exper. Appl.* **12**: 473–486.
- Francis, L. 1973. Intraspecific aggression and its effect on the distribution of *Anthopleura elegantissima* and some related anemones. *Biol. Bull.* **144**: 73–92.

- Fusetani, N., S. Matsunaga, and S. Konosu. 1981. Bioactive marine metabolites. II. Halistanol sulfate, an antimicrobial novel steroid sulfate from the marine sponge *Halichondria cf. moorei* Bergquist. *Tetrahedron Lett.* 22: 1987-1988.
- Gerhart, D. J. 1984. The staglandin A₂: an agent of chemical defense in the Caribbean spongonian *Plexaura homomalla*. *Mar. Ecol. Prog. Ser.* 19: 181-187.
- Gonzalez, A. I., D. M. Estrada, J. D. Martin, V. S. Martin, C. Perez, and R. Perez. 1984. New antimicrobial diterpenes from the sponge *Spongia officinalis*. *Tetrahedron* 40: 4109-4113.
- Green, G. 1977. Ecology of toxicity in marine sponges. *Mar. Biol.* 40: 207-215.
- Hashimoto, Y. 1979. *Marine Toxins and Other Bioactive Marine Metabolites*. Japan Scientific Society Press, Tokyo. 369 pp.
- Hollenbeak, K. H., F. J. Schmitz, and P. N. Kaul. 1976. Cardiotonic agents from marine sponges: isolation of histamine and N-methylated histamines. Pp. 282-287 in *Food and Drugs from the Sea, Proceedings 1974*. H. H. Webber and G. D. Ruggieri, eds. Marine Technology Society Press, Washington.
- Holm-Hansen, O., C. J. Lorenzen, R. W. Holmes, and J. D. H. Strickland. 1965. Fluorometric determinations of chlorophyll. *J. Cons. Perm. Internat. Explor. Mer.* 30: 3-15.
- Jackson, J. B. C. 1977. Competition on marine hard substrata: the adaptive significance of solitary and colonial strategies. *Am. Nat.* 111: 743-767.
- Jackson, J. B. C., and L. Buss. 1975. Allelopathy and spatial competition among coral reef invertebrates. *Proc. Natl. Acad. Sci. U. S. A.* 72: 5160-5163.
- Jassby, A. D., and T. Platt. 1976. Mathematical formulation of the relationship between photosynthesis and light for phytoplankton. *Limnol. Oceanogr.* 21: 540-547.
- Johannes, R. E., and W. J. Wiebe. 1970. Method for determination of coral tissue biomass and composition. *Limnol. Oceanogr.* 15: 822-824.
- Kashman, Y., A. Groweiss, R. Lidor, D. Blasberger, and S. Carmely. 1985. Latrunculins: NMR study, two toxins and a synthetic approach. *Tetrahedron* 41: 1905-1914.
- Kaul, P. N., and C. J. Sinderman, eds. 1978. *Drugs and Food from the Sea*. University of Oklahoma Press, Norman. 448 pp.
- LaBarre, S. C., and J. C. Coll. 1982. Movement in soft corals: an interaction between *Nephtea brassica* (Coelenterata: Octocorallia) and *Acropora hyacinthus* (Coelenterata: Scleractinia). *Mar. Biol.* 72: 119-124.
- LaBarre, S. C., J. C. Coll, and P. W. Sammarco. 1986a. Competitive strategies of soft corals (Coelenterata: Octocorallia): III. Spacing and aggressive interactions between alyconaceans. *Mar. Ecol. Prog. Ser.* 28: 147-156.
- LaBarre, S. C., J. C. Coll, and P. W. Sammarco. 1986b. Defensive strategies of soft corals (Coelenterata: Octocorallia) of the great Barrier reef. II. The relationship between toxicity and feeding deterrence. *Biol. Bull.* 171: 565-576.
- Lang, J. C. 1973. Interspecific aggression by scleractinian corals. 2. Why the race is not only to the swift. *Bull. Mar. Sci.* 23: 260-279.
- Manes, L. V., S. Naylor, P. Crews, and G. J. Bakus. 1985. Suvanone, a novel sesterterpene from an *Ircinia* marine sponge. *J. Org. Chem.* 50: 284-286.
- Marsh, J. A. 1970. Primary productivity of reef-building calcareous red algae. *Ecology* 51: 255-263.
- Martin, D. F., and G. M. Padilla. 1973. *Marine Pharmacognosy*. Academic Press, New York. 317 pp.
- Mayol, L., V. Piccialli, and D. Sica. 1986. Metabolites from the marine sponge *Spongimella gracilis*. *Tetrahedron* 42: 5369-5376.
- McCaffrey, E. J., and R. Endeau. 1985. Antimicrobial activity of tropical and subtropical sponges. *Mar. Biol.* 89: 1-8.
- Meinwald, J., ed. 1982. The organic chemistry of animal defense mechanisms. *Tetrahedron* 38: 1855-1970.
- Minale, L. 1978. Terpenoids from marine sponges. Pp. 175-240 in *Marine Natural Products: Chemical and Biological Perspectives, Vol. 1*. P. J. Scheuer, ed. Academic Press, New York.
- Morse, A. N. C., and D. E. Morse. 1984. Recruitment and morphosis of *Haliotis* larvae induced by molecules uniquely available at the surfaces of crustose red algae. *J. Exp. Mar. Biol. Ecol.* 75: 191-215.
- Muscatine, L., P. W. Falkowski, J. W. Porter, and Z. Dubinsky. 1984. Fate of photosynthetically-fixed carbon in light- and shade-adapted colonies of the symbiotic coral, *Stylophora pistillata*. *Proc. R. Soc. Lond. B* 22: 181-202.
- Muscatine, L., L. R. McCloskey, and R. E. Marian. 1981. Estimation of the daily contribution of carbon zooxanthellae to coral animal respiration. *Limnol. Oceanogr.* 26: 601-611.
- Nakamura, H., S. Deng, J. Kobayashi, Y. Obizumi, and Y. Hirata. 1986. Dictyoceratin-A and -B, novel antimicrobial terpenoids from Okinawan marine sponge *Hippospongia* sp. *Tetrahedron* 42: 4197-4201.
- Nakatsu, T., D. J. Faulkner, G. K. Matsumoto, and J. Clardy. 1984. Structure of the diterpene portion of a novel base from the sponge *Agelas mauritiana*. *Tetrahedron Lett.* 25: 935-9381.
- Nakatsu, T., R. P. Walker, J. E. Thompson, and D. J. Faulkner. 1983. Biologically active sterol sulfates from the marine sponge *Toxadocia zumi*. *Experientia* 39: 759-761.
- Norris, J. N., and W. Fenical. 1982. Chemical defense in tropical marine algae. *Smithson. Contrib. Mar. Sci.* 12: 417-431.
- Palumbi, S. R., and J. B. C. Jackson. 1982. Ecology of cryptic coral reef communities. II. Recovery from small disturbance events by encrusting bryozoa: the influence of "host" species and lesion size. *J. Exp. Mar. Biol. Ecol.* 64: 103-115.
- Parsons, T. R., Y. Maita, and C. M. Lalli. 1984. *A Manual of Chemical and Biological Methods for Seawater Analysis*. Pergamon Press, Oxford. 173 pp.
- Pawlik, J. R. 1986. Chemical induction of larval settlement and metamorphosis in the reef-building tube worm *Phragmatopoma californica* (Sabellariidae: Polychaeta). *Mar. Biol.* 91: 59-68.
- Porter, J. W. 1980. Primary productivity in the sea: reef corals *in situ*. Pp. 403-410 in *Primary Productivity in the Sea*, P. G. Falkowski, ed. Plenum Press, New York.
- Porter, J. W. 1985. The maritime weather of Jamaica: its effects on annual carbon budgets of the massive reef-building coral *Montastrea annularis*. *Proc. Fifth Internat. Coral Reef Congr., Tahiti* 6: 363-379.
- Porter, J. W. 1987. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (South Florida). Reef-building corals. *U. S. Fish. Wildl. Serv. Biol. Rep.* 82: 1-23.
- Porter, J. W., L. Muscatine, Z. Dubinsky, and P. G. Falkowski. 1984. Primary production and photoadaptation in light- and shade-adapted colonies of the symbiotic coral, *Stylophora pistillata*. *Proc. R. Soc. Lond. B* 216: 161-180.
- Reiswig, H. M. 1971. *In situ* pumping activities of tropical Demospongia. *Mar. Biol.* 9: 38-50.
- Richardson, C. A., P. Dustan, and J. C. Lang. 1979. Maintenance of living space by sweeper tentacles of *Montastrea cavernosa*, a Caribbean reef coral. *Mar. Biol.* 55: 181-186.
- Rosenthal, G. A., and D. H. Janzen, eds. 1979. *Herbivores: Their Interactions with Secondary Plant Metabolites*. Academic Press, New York. 718 pp.
- Russell, F. E., and P. R. Saunders, eds. 1967. *Animal Toxins*. Pergamon Press, New York. 428 pp.
- Sammarco, P. W., J. C. Coll, and S. LaBarre. 1985. Competitive strategies of soft corals (Coelenterata: Octocorallia). II. Variable de-

- fensive responses and susceptibility to scleractinian corals. *J. Exp. Mar. Biol. Ecol.* **91**: 199–216.
- Sammarco, P. W., J. C. Coll, S. LaBarre, and B. Willis. 1983. Competitive strategies of soft corals (Coelenterata: Octocorallia): Allelopathic effects on selected scleractinian corals. *Coral Reefs* **1**: 173–178.
- Scheuer, P. J., ed. 1985. The organic chemistry of marine products. *Tetrahedron* **41**: 979–1108.
- Sheppard, C. R. C. 1979. Interspecific aggression between reef corals with reference to their distribution. *Mar. Ecol. Prog. Ser.* **1**: 237–247.
- Sheppard, C. R. C. 1982. Coral populations on reef slopes and their major controls. *Mar. Ecol. Prog. Ser.* **7**: 83–115.
- Sigel, M. M., L. L. Wellham, W. Lichter, L. E. Dudek, J. L. Gargus, and A. H. Lucas. 1969. Anticellular and antitumor activity of extracts from tropical marine invertebrates. Pp. 281–294 in *Food-Drugs from the Sea Proceedings, 1969*, H. W. Youngken, ed. Marine Technology Society Press, Washington.
- Steinberg, P. D. 1984. Algal chemical defense against herbivores: allocation of phenolic compounds in the kelp *Alaria marginata*. *Science* **223**: 405–406.
- Stierle, D. B., and D. J. Faulkner. 1980. Metabolites of three marine sponges of the genus *Plakortis*. *J. Org. Chem.* **45**: 3396–3401.
- Sullivan, B., D. J. Faulkner, and L. Webb. 1983. Siphonodictidine, a metabolite of the borrowing sponge *Siphonodictyon* sp. that inhibits coral growth. *Science* **221**: 1175–1176.
- Tachibana, K., P. J. Scheuer, Y. Tsukitani, H. Kikuchi, D. van Engen, J. Clardy, Y. Gopichand, and F. Schmitz. 1981. Okadaic acid, a cytotoxic polyether from two marine sponges of the genus *Hali-chondria*. *J. Am. Chem. Soc.* **103**: 2469–2471.
- Targett, N. M., S. S. Bishop, O. J. McConnell, and J. A. Yoder. 1983. Antifouling agents against the benthic marine diatom *Nannicula salinicola*: homarine from the gorgonians *Leptogorgia virgulata* and *L. setacea* and analogs. *J. Chem. Ecol.* **9**: 817–829.
- Targett, N. M., and M. Isman, eds. 1986. Proceedings of the Second International Society of Chemical Ecology Meeting. *J. Chem. Ecol.* **12**: 949–1203.
- Targett, N. M., T. Targett, N. Vrolijk, and J. Ogden. 1986. The effect of macrophyte secondary metabolites on feeding preferences of the herbivorous parrotfish *Sparisoma radians*. *Mar. Biol.* **92**: 141–148.
- Thompson, J. E. 1985. Exudations of biologically-active metabolites in a sponge (*Aplysina fistularis*). I. Biological evidence. *Mar. Biol.* **88**: 23–26.
- Thompson, J. E., R. P. Walker, D. J. Faulkner. 1985. Screening and bioassays for biologically active substances from forty marine sponge species from San Diego, Ca., U. S. A. *Mar. Biol.* **88**: 11–22.
- Thompson, J. E., R. P. Walker, S. J. Wratten, and D. J. Faulkner. 1982. A chemical defense mechanism for the nudibranch *Cadlina luteomarginata*. *Tetrahedron* **38**: 1865–1873.
- Tursch, B. 1976. Some recent developments in the chemistry of alcyonaceans. *Pure Appl. Chem.* **48**: 1–6.
- Tursch, B., J. C. Braekman, D. Dalozze, and M. Kaisin. 1978. Terpenoids from coelenterates. Pp. 247–296 in *Marine Natural Products: Chemical and Biological Perspectives Vol. II*, P. J. Scheuer, ed. Academic Press, New York.
- Tursch, B., and A. Tursch. 1982. The soft coral community on a sheltered reef quadrat at Laing Island (Papua New Guinea). *Mar. Biol.* **68**: 321–332.
- Wahle, C. E. 1980. Detection, pursuit and overgrowth of tropical gorgonians by milleporid hydrocorals: Perseus and Medusa revisited. *Science* **209**: 689–691.
- Walker, R. P., J. E. Thompson, and D. J. Faulkner. 1985. Exudation of biologically active metabolites in a sponge *Aplysina fistularis*. II. Chemical evidence. *Mar. Biol.* **88**: 27–32.
- Webb, L., and J. C. Coll. 1983. Effects of alcyonarian coral terpenes on scleractinian coral photosynthesis and respiration. *Toxicon Suppl.* **3**: 485–488.
- Wellington, G. M. 1980. Reversal of digestive interactions between Pacific reef corals: mediation by sweeper tentacles. *Oecologia* **47**: 340–343.
- Whittaker, R. H., and P. P. Feeny. 1971. Allelochemicals: chemical interactions between species. *Science* **171**: 757–770.